



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Halsted's book is lucidly written, the translation is at once faithful and smooth."

We congratulate Professor Halsted on the well-merited success that his book has attained. Sooner or later it will no doubt exercise a profound influence on the teaching of geometry in our country.

---

## A METHOD FOR THE SOLUTION OF SIMULTANEOUS QUADRATIC EQUATIONS OF THE SYMMETRIC TYPE.

---

By B. E. MITCHELL, Nashville, Tennessee.

---

The following method is suggested but not developed in Fine's *College Algebra*.

The most general quadratic of the symmetric type between two variables is  $a(x^2+y^2)+2bxy+c(x+y)+d=0$ .

Let  $x$  and  $y$  be the roots of  $u^2+pu+q=0$ . Then

$$u = \begin{cases} x = \frac{1}{2} \left[ -p \pm \sqrt{p^2 - 4q} \right] \\ y = \frac{1}{2} \left[ -p \mp \sqrt{p^2 - 4q} \right] \end{cases} \dots (I).$$

The double sign is used in each because they are the roots of a symmetric equation.

From the theory of quadratic equations we have

$$x+y=-p\dots(1),$$

and

$$xy=q\dots(2),$$

from which we calculate

$$x^2+y^2=p^2-2q\dots(3).$$

These are the three symmetric forms that occur in the general equation above, and are the only ones of the second degree.

Let us solve two simultaneous equations by this method.

$$\begin{cases} a_1(x^2+y^2)+2b_1xy+c_1(x+y)+d_1=0\dots(1). \\ a_2(x^2+y^2)+2b_2xy+c_2(x+y)+d_2=0\dots(2). \end{cases}$$

Then

$$\begin{cases} a_1p^2-2(a_1-b_1)q-c_1p+d_1=0\dots(3). \\ a_2p^2-2(a_2-b_2)q-c_2p+d_2=0\dots(4). \end{cases}$$

Each of the equations (3) and (4) is linear in  $q$ . Solving them for  $q$ , we have

$$2q = \frac{a_1 p^2 - c_1 p + d_1}{a_1 - b_1} = \frac{a_2 p^2 - c_2 p + d_2}{a_2 - b_2} \dots (5).$$

Considering equation (5) we have the following:

The coefficient of  $p^2$  is  $a_2 b_1 - a_1 b_2 = - \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix}$  which we call  $A$ .

The coefficient of  $p$  is  $a_1 c_1 - a_2 c_2 - b_1 c_2 + b_2 c_1 = - \begin{vmatrix} a_1 & a_2 & 1 \\ b_1 & b_2 & 1 \\ c_1 & c_2 & 0 \end{vmatrix}$  which we call  $B$ .

The constant term is  $a_2 d_1 - b_2 d_1 - a_1 d_2 + b_1 d_2 = - \begin{vmatrix} a_1 & a_2 & 1 \\ b_1 & b_2 & 1 \\ d_1 & d_2 & 0 \end{vmatrix}$  which we call  $C$ .

Hence we have  $Ap^2 + Bp + C = 0$  yielding generally two values for  $p$ , and from (5) we may calculate the two corresponding values of  $q$ . We have then the two equations in  $u$ :

$$\begin{aligned} u^2 + p_1 u + q_1 &= 0, \\ u^2 + p_2 u + q_2 &= 0, \end{aligned}$$

which give four pairs of values for  $x$  and  $y$  when substituted in (I).

In the application of the method to numerical problems it will usually be found easier to use it as a process rather than the above equations as formulae.

Let us apply the method to three numerical problems taken from Fine's *College Algebra*.

$$(1) \begin{cases} x + y = 5. \\ xy + 36 = 0. \end{cases}$$

Hence  $p = -5$  and  $q = -36$ , so  $u^2 - 5u - 36 = 0$ , and  $u = \begin{cases} x = 9 \text{ and } -4. \\ y = -4 \text{ and } 9. \end{cases}$

$$(2) \begin{cases} x^2 + xy + y^2 = 21. \\ x + \sqrt{xy} + y = 7. \end{cases} \quad \begin{cases} p^2 - 2q + q = 21. \\ -p + \sqrt{q} = 7. \end{cases} \quad \begin{cases} p^2 - q = 21. \\ p^2 - q + 14p = -49. \end{cases}$$

$$p = -5, \quad q = 4. \quad u^2 - 5u + 4 = 0. \quad (u - 4)(u - 1) = 0.$$

$$u = \begin{cases} x = 1 \text{ and } 4. \\ y = 4 \text{ and } 1. \end{cases}$$

$$(3) \begin{cases} x^2 + 3xy + y^2 + 2x + 2y = 8. \\ 2x^2 + 2y^2 + 3x + 3y = 14. \end{cases}$$

Then  $p^2 - 2p + q = 8$ , and  $2p^2 - 3p - 2q = 14$ .

Eliminating  $q$  by addition,  $4p^2 - 7p - 30 = 0$ ,  $p = -2$  and  $\frac{1}{4}$ , then  $q = -0$  and  $\frac{2}{16}$ . So  $u^2 - 2u = 0 \dots (a)$ , and  $u^2 + \frac{1}{4}u + \frac{2}{16} = 0 \dots (b)$ .

$$\text{From (a), } u = \begin{cases} x = 0 \text{ and } 2. \\ y = 2 \text{ and } 0. \end{cases}$$

$$\text{From (b), } u = \begin{cases} x = \frac{1}{2} \left[ -\frac{1}{4} \pm \sqrt{\left(\frac{1}{16}\right)} \right]. \\ y = \frac{1}{8} \left[ -15 \mp \sqrt{(133)} \right]. \end{cases}$$

This method may also be used where the equations are symmetric with respect to  $x$  and  $-y$ . In this case we have

$$x-y=-p, \quad xy=-q, \quad \text{and} \quad x^2+y^2=p^2-2q.$$

By calculating other symmetric functions the method can be used successfully in solving many equations of higher degree than the second.

## SOME CONSTRUCTIONS LEADING TO CONICS.

By F. H. HODGE, Franklin College, Indiana.

Among the courses that find a place in collegiate mathematics one is usually given which involves the treatment of plane curves. Certain loci are




Fig. 1.

The four constructions which follow lead to conics. These are so simple and so directly analogous to a well known construction for the ellipse that it is scarcely possible that they are new though they are not found in current texts. The first two are described in detail, and the others merely suggested.

(1) Given two circles tangent internally at  $B$  and having the radius of the larger equal to the diameter of the smaller. Take the center of the larger circle as origin and the tangent to the smaller circle at that point as the  $x$ -axis, the  $y$ -axis being perpendicular to the  $x$ -axis. Draw a secant line through the origin, meeting the smaller circle at  $R$  and the larger circle at  $S$ . Through  $R$  draw a line parallel to the  $x$ -axis, and through  $S$  draw a line parallel to the  $y$ -axis. Call the point  $P$  in which these two lines meet. Required to find the locus of  $P$  as the secant line revolves about the origin as an axis. See Fig. 1.




Fig. 2.